

General Approach for Deriving Reference Distribution Functions for Systems out of Equilibrium by Statistical Thermodynamics

Giorgio SONNINO,^{1*} and Alessandro CARDINALI², Mustapha TLIDI¹,
Philippe PEETERS¹, György STEINBRECHER³, Alexander MILOVANOV²

¹Université Libre de Bruxelles (U.L.B.), Department of Theoretical Physics and
Mathematics, Campus de la Plaine C.P. 231 - Bvd du Triomphe, 1050 Brussels - Belgium

²EURATOM-ENEA Fusion Association, Via E.Fermi
45, C.P. 65 - 00044 Frascati (Rome) - Italy

³EURATOM-MEdC Fusion Association, University of Craiova, Faculty
of Exact Sciences, Str.A.I.Cuza Street 13, Craiova-200585, Romania

Abstract

A general approach for deriving the expression of *reference (density of) distribution functions*, $\mathcal{F}^0$, by statistical thermodynamics and the definition of local equilibrium conditions is illustrated. This procedure may be adopted for a system subject to an arbitrary number of thermodynamic forces. For concreteness, we analyze the case of a system submitted to three independent thermodynamic forces and the local equilibrium corresponds to the configuration of *minimum entropy production condition* and the *maximum entropy principle*. In this limit case, we show that the derived expression of distribution function is more general than that one, which is currently used for fitting the numerical steady-state solution obtained by simulating the Ion Cyclotron Radiation Heating (ICRH) FAST-plasmas and for describing various scenarios of tokamak plasmas. Through kinetic theory, we fixed the free parameters linking them with the external power supplies. The singularity at low energy in the proposed distribution function is related to the intermittency in the turbulent plasma. As a matter of fact, this work is not restricted to, but applied to, tokamak plasmas. Tokamak-plasmas are taken as an example of close thermodynamic systems. An application to a simple model of fully ionized tokamak-plasmas submitted to an external Ohmic source is discussed.

PACS Numbers: 52.25.Dg, 05.70.Ln, 05.20.Dd, 52.55.Fa.

Keywords: Statistical Thermodynamics, Thermodynamics of Irreversible Processes, Kinetic Theory, Tokamak-Plasmas.

*Electronic address: gsonnino@ulb.ac.be

I. INTRODUCTION

Statistical thermodynamics constitutes a powerful tool for deriving the reference density distribution functions (DDF), $\mathcal{F}^0$ (see, for instance, [1]). The DDF is obtained by starting from the Einstein formula linking the (density of) probability of a fluctuation, $\mathcal{F}$, with the entropy variation, ΔS , associated with the fluctuations from the state of equilibrium [2], [3] and [4]

$$\mathcal{F} = \mathcal{N}_0 \exp[\Delta S] \quad (1)$$

In Eq. (1), Boltzmann's constant is set equal to 1 and $\mathcal{N}_0$ is a normalization constant that ensures the sum of all probabilities equals one. Notice that, since ΔS is the change of entropy starting from equilibrium, we have $\Delta S < 0$. Prigogine generalized Eq. (1), which applies only to adiabatic or isothermal transformations, by introducing the entropy production due to fluctuations. Let us consider a system characterized by n variables $A_1, A_2, \dots, A_n$. The (local) equilibrium values are $A_1^0, A_2^0, \dots, A_n^0$. Denoting by $\alpha_i = A_i - A_i^0$ ($i = 1 \dots n$) the n deviations of the thermodynamic quantities from their equilibrium value, Prigogine proposed that the probability distribution of finding a state in which the values α_i lie between α_i and $\alpha_i + d\alpha_i$ is given by [5]

$$\mathcal{F} = \mathcal{N}_0 \exp[-\Delta_I S] \quad (2)$$

where we have introduced the *dimensionless (density of) entropy production* $\Delta_I S$ [6]

$$\Delta_I S = \frac{\tau}{n} \sigma \geq 0 \quad (3)$$

Here τ , n and σ denote the *relaxation time*, the *number density* and the *entropy source strength* (per unit volume), respectively. The negative sign in Eq. (2) is due to the fact that, during the processes, $-\Delta_I S \leq 0$. Indeed, if $-\Delta_I S$ were positive, the transformation $\alpha_i \rightarrow \alpha'_i$ would be a spontaneous irreversible change and thus be incompatible with the assumption that the initial state is a (local) equilibrium state [5]. Using this notation, the thermodynamic forces X^κ and the thermodynamic fluxes J_κ are

$$X^\kappa = \frac{\partial \Delta_I S}{\partial \alpha_\kappa} \quad ; \quad J_\kappa = \frac{d\alpha_\kappa}{dt} \quad (4)$$

and the expressions for $d_I S$ and σ read [7]

$$\begin{aligned} d_I S &= \sum_{\kappa=1}^n X^\kappa d\alpha_k \quad \text{or} \quad \frac{d_I S}{dt} = \sum_{\kappa=1}^n J_\kappa X^\kappa \geq 0 \\ \frac{d_I S}{dt} &= \int_{\Omega} \sigma d\mathbf{x} \end{aligned} \quad (5)$$

where $d\mathbf{x}$ is a (spatial) volume element, and the integration is over the entire space Ω occupied by the plasma. Note that the probability density function (2) remains unaltered for flux-force transformations, $X_\kappa \rightarrow X'_\kappa$ and $J_\kappa \rightarrow J'_\kappa$, leaving invariant the entropy production [8]

$$\frac{d_I S}{dt} = \sum_{\kappa=1}^n J_\kappa X^\kappa = \sum_{\kappa=1}^n J'_\kappa X'^\kappa \quad (6)$$

The expressions for the reference distribution function, $\mathcal{F}^0$, can now be obtained by imposing a set of minimum conditions able to identify, uniquely, the local equilibrium. These conditions should be established according to the particular physical situation that we are analyzing. It is important to stress that the set of restrictions, imposed by very general physical principles, including scale invariance, gives automatically, for some range of parameters, a distribution function whose singularity can be interpreted in the terms of intermittency on turbulent plasma (see Appendix). This procedure is quite general and it can be applied to a system subject to an arbitrary number of thermodynamic forces. However for concreteness, we limit ourselves to study the particular case of a system submitted to only three thermodynamic forces. The local equilibrium condition is determined by imposing the *minimum entropy production condition* and the *maximum entropy principle*.

The main objective of this paper is to derive the expression of $\mathcal{F}^0$ for the case of a system subject to three independent thermodynamic forces by statistical thermodynamics. The link between the free parameters with the external power supply and the entropy source strength of the confined tokamak-plasma is performed by kinetic theory. This example of calculation illustrates how this procedure applies to the general case of systems subject to a larger number of thermodynamic forces. We mention that the present state of the art in tokamak modelling allows obtaining distribution functions from sophisticated numerical codes (e.g., NUBEAM, ASCOT, PENCIL for NBI, and TORIC, PION for ICRH). These numerical codes can actually be used for calculating numerically the particle distribution function for different power injection schemes and different levels of approximations in treating sources, collision operators and particle motion in the equilibrium fields. The scope of the present

work is to provide a reasonably complete class of model distribution functions to be used in gyrokinetic or hybrid gyrokinetic-fluid simulation of nonlinear dynamics of drift Alfvén waves in burning plasmas of fusion interest. The coupling of such gyrokinetic or hybrid gyrokinetic-fluid simulation with power deposition codes, like those mentioned above, is a challenging task, which is underway. In the meantime, sufficiently accurate description of model particle distribution functions are needed, which nowadays are typically chosen as "reasonable" model functions of the particle constants of motion (often just a Maxwellian, as in GTC and GYRO, or more generally an anisotropic Maxwellian or slowing down for the case of HMGC, M3D and NOVA). In this work, the constraints that non-equilibrium statistical mechanics impose on the adopted model distribution functions are derived and a class of model distribution function is proposed, whose usefulness is therefore, the readiness to be adopted in gyrokinetic or hybrid gyrokinetic-fluid simulation of nonlinear dynamics of drift Alfvén waves in burning plasmas of fusion interest. At the present time, none of the above-mentioned codes can be used in the same way, for the intrinsic difficulty of an actual integrated simulation. The advantage of the model distribution functions obtained in this work is therefore evident. As a matter of fact the work is not restricted to but applied to tokamak plasmas. The fundamental issue here is that the detailed application to tokamaks comes in only when the specific form of constants of motion in the equilibrium fields is adopted. Before that point, there is a construction of the particle distribution function out of the equilibrium, as we expect that to be in the case of a tokamak. The final part of the work is addressed to study the inverse problem: given $\mathcal{F}^0$, we are looking for the Fokker-Planck equation admitting $\mathcal{F}^0$ as the steady-state solution.

The paper is organized as follows. In Section (II) we derive the general expression of the reference density of distribution probability, $\mathcal{F}^0$, by a purely thermodynamic approach. The parameters, entering in the expression of $\mathcal{F}^0$, are determined by kinetic theory in Section III by adopting a model for the tokamak-plasma and the external sources. Section IV addresses the following questions :

- *For collisional tokamak-plasmas, how much is the deviation of the reference DDF, $\mathcal{F}^0$, from the Maxwellian ?*
- *Does this deviation coincide with the one estimated by the neoclassical theory (see, for example, Ref. [12]) ?*

These are crucial questions because, when a new approach is proposed, one of the main tasks is to show that the new approach is also able to recover the results, which are well established in literature. In our case, we should ensure that the expression that we found for the reference DDF, $\mathcal{F}^0$, *coincides exactly* with the one predicted by the neoclassical theory for collisional tokamak-plasmas. As we shall see in this question, the answer to this question is positive. Some concluding remarks can be found in the Section V.

II. THERMODYNAMIC DERIVATION OF THE DISTRIBUTION FUNCTION. GENERAL CONSIDERATIONS.

In this section, we derive the form of $\mathcal{F}^0$, the *reference density of probability function*, by a pure thermodynamic approach. We limit ourselves to study a system subject to three thermodynamic forces. As usual, the gyrokinetic (GK) theory makes often use of an initial distribution function of guiding centers. In the GK simulations, as well as in the GK theory, this initial distribution function is usually taken as a reference DDF if it depends only on the invariants of motion and it evolves slowly from the local equilibrium state i.e., in such a way that the guiding centers remain confined for sufficiently long time. After a short transition time, the state of the plasma remains close to the *reference state*, which results to be a small deviation of the (local) equilibrium state. The expression of the coefficients of the $\mathcal{F}^0$ will be determined in the next section by kinetic theory. The reference DDF is obtained by perturbing the local equilibrium state. To determine the reference DDF we shall follow the following strategy. First we consider open thermodynamic systems obeying to Prigogine statistical thermodynamics. Successively, we define the local equilibrium state by adopting a minimal number of hypotheses. Finally, we link the Prigogine probability distribution function with particles DDF. The expression of the *local equilibrium state* is fixed by adopting a set of minimal number of *ad hoc* assumptions. We restrict ourselves by analyzing the case of an axisymmetric magnetically confined plasma. After having performed the guiding center transformation, the necessary variables for describing the system reduce to four independent variables [6]. In addition, we consider the case where the system is subject to three thermodynamic forces, associated to the three fluctuations α_1 , α_2 and α_3 with α_3 identified as w , the particle kinetic energy per unit mass. Hence, $w = (v_{\parallel}^2 + v_{\perp}^2)/2$ where $v_{\parallel}$ denotes the parallel component of particle's velocity, which may actually be parallel or

antiparallel to the magnetic field and $v_\perp$ is the absolute value of the perpendicular velocity. Now we should define the local equilibrium state. We assume that (see also the footnote [31])

The equilibrium state corresponds to the values $\alpha_1 = 0$ and $\alpha_2 = 0$ for which the entropy production tends to reach an extreme.

Hence, α_1 and α_2 are two Prigogine's fluctuations [for their exact definition see Ref. [5] and [13]. For fully ionized collisional plasma, these fluctuations are linked to the thermodynamic forces by Eq. (59) (see the Section IV)]. At this state, the deviation of $\Delta_I S$ from its equilibrium value is given by [5] and [13]

$$-\Delta_I S = g_0(w) - \frac{1}{2} \sum_{i,j=1}^2 g_{ij}(w) \alpha_i \alpha_j + h.o.t. \quad \text{with} \quad (7)$$

$$g_0(w) \equiv -\Delta_I S |_{\alpha_1=\alpha_2=0}$$

where *h.o.t.* stands for *higher order terms*. Therefore, the general DDF becomes $\mathcal{F}^0$ when the expression of entropy production is given by Eq. (7). Coefficients g_{ij} denote the second derivatives of $\Delta_I S$ with respect to the α_i variables. With these coefficients we may form a positive definite matrix. This matrix can be diagonalized by obtaining

$$-\Delta_I S = g_0(w) - \sum_{i,j=1}^2 \tilde{c}_i(w) \delta_{ij} \xi_i^2 + h.o.t. \quad (8)$$

where $\tilde{c}_i(w) \delta_{ij}$ is a positive definite matrix and δ_{ij} denotes Kronecker's delta. Eq. (8) allows describing the entire process in terms of two *independent* processes linked to the two independent fluctuations ξ_1 and ξ_2 . As mentioned, $\mathcal{F}^0$ is determined by four variables. In order to simplify, as much as possible, the expression of the reference density of probability function, we shall now introduce a set of convenient variables. We introduce two variables, ζ_1 and ζ_2 , of the type, *degrees of advancement* (for a rigorous definition, see any standard textbook such as, for example, Ref. [13]. See also the footnote [32]), and the remaining variable, ζ_3 , is chosen such that the fluctuations ξ_1 and ξ_2 depend on the variables ζ_1 and ζ_2 , but not on variable ζ_3 . The reference density of probability function will be then determined by the following set of coordinates: $(w, \zeta_1, \zeta_2, \zeta_3)$. We recall that the remaining two coordinates: *the toroidal angle*, ϕ , and *the gyro-phase angle*, Φ , are unnecessary due to the axisymmetric configuration and the guiding center transformation. ζ_1 and ζ_2 are associated to the two independent fluctuations ξ_1 and ξ_2 (therefore, they are also associated to the two

independent thermodynamic forces). We may expand the fluctuations ξ_1 and ξ_2 in Taylor's series of variables (ζ_1, ζ_2) by obtaining

$$\begin{aligned}\xi_1 &= \left(\frac{\partial \xi_1}{\partial \zeta_1} \right)_{\zeta_1^0, \zeta_2^0} (\zeta_1 - \zeta_1^0) + h.o.t. \\ \xi_2 &= \left(\frac{\partial \xi_2}{\partial \zeta_2} \right)_{\zeta_1^0, \zeta_2^0} (\zeta_2 - \zeta_2^0) + h.o.t.\end{aligned}\quad (9)$$

In Eqs. (9) we have taken into account that ζ_1 and ζ_2 are two degrees of advancement variables and that ξ_1 and ξ_2 are two independent fluctuations (see the footnote [33]). Hence, the reference distribution function (2) takes the form

$$\begin{aligned}d\hat{\mathcal{F}}^0 &= \mathcal{P}(w) \exp[-c_1(w)(\zeta_1 - \zeta_1^0)^2] \exp[-c_2(w)(\zeta_2 - \zeta_2^0)^2] d\Gamma \quad \text{with} \\ \mathcal{P}(w) &\equiv \exp[g_0(w)] = \exp[-\Delta_I S]_{\xi_1=\xi_2=0} \quad \text{and} \quad c_i(w) \equiv \tilde{c}_i(w) \left(\frac{\partial \xi_i}{\partial \zeta_i} \right)_{\zeta_1^0, \zeta_2^0}^2 \quad (i = 1, 2)\end{aligned}\quad (10)$$

The phase space volume element $d\Gamma = d\mathbf{x}d\mathbf{v}$ is linked to the new volume element $d\hat{\Gamma} = dw d\zeta_1 d\zeta_2 d\zeta_3 d\phi d\Phi$ by

$$d\Gamma = |\mathcal{J}| d\hat{\Gamma} \quad (11)$$

where $|\mathcal{J}|$ is the Jacobian between $d\Gamma$ and $d\hat{\Gamma}$. To proceed further in our analysis, we should complete our definition of *equilibrium state*. For this, we may think to w as a random variable satisfying two restrictions

$$\begin{aligned}\mathbb{E}[w] &\sim T/m > 0 \\ \mathbb{E}[\ln(w)] &\sim \Psi(T/m)\end{aligned}\quad (12)$$

where T is temperature and m denotes particle's mass. $\mathbb{E}[\]$ and $\Psi(x)$ are the expectation operation and the digamma function, respectively. The equilibrium state is fully determined by assuming the validity of the following *Principle of Maximum Entropy*.

In the phase space Γ , at the externalizing values $\alpha_1 = \alpha_2 = 0$ (or, which is the same, at $\xi_1 = \xi_2 = 0$), the system tends to evolve towards maximal entropy configurations under the constraints given by Eqs (12).

Under this assumption, the random variable w is modeled by a gamma distribution [14]

$$\mathcal{P}(w/\Theta) = \mathcal{N}_0 \left(\frac{w}{\Theta} \right)^{\gamma-1} \exp[-w/\Theta] \quad (13)$$

where we have introduced the *scale parameter* Θ and the *shape parameter* γ . Conditions (12) fix the parameters of the Gamma distribution (see, for example, Ref. [15]). The motivation

for the choice of the restrictions given by Eqs(12), as well as the special mathematical properties of the resulting PDF, are given in the Appendix. Hence, the maximum entropy principle allows writing the reference state, Eq. (10), as

$$d\hat{\mathcal{F}}^0 = \mathcal{N}_0 \left(\frac{w}{\Theta} \right)^{\gamma-1} \exp[-w/\Theta] \exp[-c_1(w/\Theta)(\zeta_1 - \zeta_1^0)^2] \exp[-c_2(w/\Theta)(\zeta_2 - \zeta_2^0)^2] | \mathcal{J} | d\hat{\Gamma} \quad (14)$$

If we interpret our reference DDF as a time and ensemble average of the physical DDF describing turbulent plasma, then the singularity at $w = 0$ for $0 < \gamma < 1$ can be related to the intermittency (see Appendix). Notice that at the point with coordinates $(\alpha_1, \alpha_2, \alpha_3) = [0, 0, (\gamma - 1)\Theta]$, the system satisfies the principle of maximum entropy and the entropy production reaches its extreme value.

Let us now consider the coefficients $c_1(w/\Theta)$ and $c_2(w/\Theta)$. As an example of a closed thermodynamic system (i.e., the system can exchange energy but *not* matter with an outside system), we analyze collisional tokamak-plasmas where the magnetic configuration can be brought into the form (see, for instance, Ref. [12])

$$\mathbf{B} = -\frac{1}{2\pi} \nabla\psi \times \nabla\phi + F(\psi) \nabla\phi \quad (15)$$

The particle energy is $\mathcal{E} = mw + qA_E \sim mw$, with m , q and A_E denoting the *mass particle*, the *charge particle* and the *electric potential*, respectively. Moreover, ψ indicates the *poloidal magnetic flux* and F is the *characteristic of axisymmetric toroidal field* depending on the *surface function* ψ . We identify ζ_1 , ζ_2 and ζ_3 with the *toroidal angular moment*, P_ϕ , the variable, λ , and ψ , respectively. These quantities are defined as (for a rigorous definition, see any standard textbook such as, for example, [16])

$$P_\phi = \psi + \frac{B_0}{\Omega_{0c}} \frac{F v_\parallel}{|B|} \quad (16)$$

$$\lambda = \frac{\sin^2 \theta_P}{2 |B|} = \frac{\mu}{w} \quad \text{with} \quad \mu = \frac{v_\perp^2}{2 |B|}$$

Here Ω_{0c} is the *cyclotron frequency* associated with the magnetic field along the magnetic axis, B_0 . $|B|$ and θ_P denote the magnetic field intensity and the *pitch angle*, respectively. We expand these coefficients in terms of w . Considering that w/Θ behaves as an entropy, and that $(P_\phi - P_{\phi 0})^2$ behaves as $v_\parallel^2$, for coherence with Eqs (1) and (2), the expansions should be truncated at the zero order for c_1 and at the first order for c_2 (ζ_1^0 and ζ_2^0 are evaluated

at the zero order). Hence

$$\begin{aligned} c_1(w/\Theta) &= c_1^{(0)} + h.o.t. \\ c_2(w/\Theta) &= c_2^{(0)} + c_2^{(1)} \frac{w}{\Theta} + h.o.t. \end{aligned} \quad (17)$$

The expressions for c_1 and c_2 can be brought in the more convenient form

$$\begin{aligned} c_1(w/\Theta) &= c_1^{(0)} \equiv \left(\frac{1}{\Delta P_\phi} \right)^2 = const. \\ c_2(w/\Theta) &= \frac{1}{\Delta \lambda_0} \left(\frac{\Delta \lambda_0}{\Delta \lambda_1} + \frac{w}{\Theta} \right) \geq 0 \end{aligned} \quad (18)$$

Finally, the expression for the (density of) probability distribution function $\mathcal{F}^0$ reads

$$\mathcal{F}^0 = \mathcal{N}_0 \left(\frac{w}{\Theta} \right)^{\gamma-1} \exp[-w/\Theta] \exp \left[- \left(\frac{P_\phi - P_{\phi 0}}{\Delta P_\phi} \right)^2 \right] \exp \left[- \left(\frac{\Delta \lambda_0}{\Delta \lambda_1} + \frac{w}{\Theta} \right) \frac{(\lambda - \lambda_0)^2}{\Delta \lambda_0} \right] | \mathcal{J} | \quad (19)$$

where $\mathcal{N}_0$ ensures normalization to unity

$$\int_{\hat{\Omega}} d\hat{\mathcal{F}}^0 = \int_{\hat{\Omega}} \mathcal{F}^0 d\hat{\Gamma} = 1 \quad (20)$$

Observe that the presence of the free parameter $c_2^{(0)}$ is crucial. Indeed, as we shall show in the Section IV [in particular, see Eq. (81)], the absence of $c_2^{(0)}$ precludes the possibility of identifying the DDF, given by Eq. (19), with the one estimated by the neoclassical theory for collisional tokamak-plasmas (see, for example, Ref. [12]). In addition, it allows describing more complex physical scenarios such as, for example, the *modified bi Maxwellian distribution function* [17]. Last and not least, in some physical circumstances, the presence of $c_2^{(0)}$ is essential to ensure the normalization of the DDF. Thermodynamics has been able to determine the shape of the DDF, but it is unable to fix the seven parameters $\Theta, \gamma, P_{\phi 0}, \lambda_0, \Delta P_\phi, \Delta \lambda_0, \Delta \lambda_1$. These coefficients have to be calculated in the usual way by kinetic theory. Figs (1)-(2), (3)-(4), and (5)-(6) illustrate three plots of Eq. (19) (estimated for unit values of the Jacobian and the normalization coefficient) corresponding to the values $w = E\Theta$, $\gamma = 1 + E$ (with E denoting the Euler number), $P_\phi = P_{\phi 0}$ and $\lambda = \lambda_0$, respectively.

III. DETERMINATION OF THE PARAMETERS

This section is devoted to the determination of the value of the parameters appearing in Eq. (19) for $\mathcal{F}^0$. As an example of application, we analyze tokamak-plasmas in collisional transport regimes. To accomplish this task we should




FIG. 1: Distribution function, Eq. (19), (per unit values of the Jacobian and the normalization factor) computed at $\gamma = 1 + E$ (E indicates the Euler number), $w = E\Theta$, $\Delta P_\phi = 22.360$, $\Delta\lambda_0 = 50.00$ and $\Delta\lambda_1 = 30.2031$.




FIG. 2: Contour plot of Fig. 1.

- Adopt a model for tokamak-plasmas;
- Model the source terms.

Calculations should be performed by a kinetic approach. For definiteness, the particle density n_α and the average velocity $\mathbf{u}_\alpha$ of each species α (with $\alpha = e$ for electrons and $\alpha = i$ for ions) satisfy the equations

$$n_\alpha(\mathbf{x}) = \int_{\mathcal{V}} d\mathbf{v} \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \quad (21)$$

$$n_\alpha(\mathbf{x}) \mathbf{u}^\alpha(\mathbf{x}) = \int_{\mathcal{V}} d\mathbf{v} \mathbf{v} \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \quad \text{with} \quad \alpha = (e, i) \quad (22)$$

where $\mathcal{V}$ is the velocity-volume in the phase-space. In addition, we should construct representation (19) in such a way that temperature T_α does coincide with the exact value of the temperature of each species. This implies

$$n_\alpha(\mathbf{x}) T_\alpha(\mathbf{x}) = \frac{1}{2} m_\alpha \int_{\mathcal{V}} d\mathbf{v} |\mathbf{v} - \mathbf{u}_\alpha|^2 \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \quad \text{with} \quad \alpha = (e, i) \quad (23)$$




FIG. 3: Distribution function, Eq. (19), (per unit values of the Jacobian and the normalization factor) computed at $\gamma = 1 + E$, $\Delta P_\phi = 22.360$ and $\lambda = \lambda_0$.




FIG. 4: Contour plot of Fig. 3.




FIG. 5: Distribution function, Eq. (19), (with the Jacobian and the normalization factor set to 1) computed at $\gamma = 1 + E$, $P_\phi = P_{\phi 0}$, $\Delta\lambda_0 = 11.111$ and $\Delta\lambda_1 = 50.00$.




FIG. 6: Contour plot of Fig. 5.

where m_α is the mass particle of specie α . To the Eqs (21)-(23), we should add the equations warranting the validity of the energy and entropy balances. To this end, we consider three kind of external energetic sources: the total Ohmic heat, $\Delta\mathcal{E}_{Oh.}^\alpha$, supplied from outside, the energy $\Delta\mathcal{E}_{NBI}^\alpha$, supplied by neutral beam injection (NBI), and the energy of the minority population heated by ion cyclotron resonant heating (ICRH) $\Delta\mathcal{E}_{ICRH}^\alpha$. Here, we shall not deal with burning fusion plasmas and the loss due to the Bremsstrahlung effect is neglected. At the end, the energy balance equation reads

$$\frac{1}{2}m_\alpha \int_{\{\Omega\mathcal{V}\}} d\mathbf{\Gamma} |\mathbf{v} - \mathbf{u}_\alpha|^2 \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) = \Delta\mathcal{E}_{Oh.}^\alpha + \Delta\mathcal{E}_{NBI}^\alpha + \Delta\mathcal{E}_{ICRH}^\alpha - \Delta\mathcal{E}_L^\alpha \quad (24)$$

where $\Delta\mathcal{E}_L^\alpha$ is the energy loss and we have neglected the electric contribution with respect to the kinetic energy. Let us now consider the entropy balance. The total entropy, s , is defined as [6]

$$n_\alpha(\mathbf{x})s_\alpha(\mathbf{x}) = - \int_{\mathcal{V}} d\mathbf{v} \mathcal{F}^0(\mathbf{v}, \mathbf{x}) \ln\left(\frac{h^3}{em_\alpha^3} \mathcal{F}^0(\mathbf{v})\right) = - \int_{\mathcal{V}} d\mathbf{v} \mathcal{F}^0(\mathbf{v}, \mathbf{x}) \ln \mathcal{F}^0(\mathbf{v}) \quad (25)$$

Here h is the Planck constant, e is the (positive) magnitude of the electric charge and m_α the mass of the species α . This equation is, however, inconvenient for setting the values of the parameters, because it involves the total entropy density that, at this stage, is not a known quantity. It is more useful to study separately the two entropy contributions, $d_E S^\alpha$ and $d_I S^\alpha$, of the total entropy S^α :

$$dS^\alpha = d_E S^\alpha + d_I S^\alpha \quad \text{with} \quad S^\alpha = \int_{\Omega} d\mathbf{x} n_\alpha(\mathbf{x})s_\alpha(\mathbf{x}) \quad (26)$$

$d_E S^\alpha$ represents the amounts of entropy crossing the boundaries. The total entropy flux of species α is given by two contributions: the *convective entropy flux*, $n_\alpha s_\alpha \mathbf{u}^\alpha$ and the *conductive entropy flux*, $\mathbf{J}_S^\alpha$. On the contrary, $d_I S^\alpha$ represents the source entropy, due to internal processes, which remains within the system. In terms of the conductive entropy flux, $\mathbf{J}_S^\alpha$, and the entropy source strength, σ^α , Eq. (26) takes the form

$$\frac{dS^\alpha}{dt} = - \int_{\Sigma} d\mathbf{A} \cdot [n_\alpha s_\alpha \mathbf{u}^\alpha + \mathbf{J}_S^\alpha(\mathbf{x})] + \int_{\Omega} d\mathbf{x} \sigma^\alpha(\mathbf{x}) \quad (27)$$

where Σ denotes the boundary. According to the kinetic theory we have [7]

$$\begin{aligned} \mathbf{J}_S^\alpha(\mathbf{x}) &= - \int_{\mathcal{V}} d\mathbf{v} [\mathbf{v} - \mathbf{u}^\alpha(\mathbf{x})] \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \ln\left(\frac{h^3}{em_\alpha^3} \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x})\right) \\ &= - \int_{\mathcal{V}} d\mathbf{v} [\mathbf{v} - \mathbf{u}^\alpha(\mathbf{x})] \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \ln \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \end{aligned} \quad (28)$$

and

$$\sigma^\alpha = \frac{n_\alpha}{\tau_\alpha} \Delta_I S^\alpha = - \sum_{\beta=e,i} \int_{\mathcal{V}} d\mathbf{v} [\ln \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x})] \mathcal{K}^{\alpha\beta} \quad (29)$$

where $\mathcal{K}^{\alpha\beta}$ denotes the collisional operator of species α due to β and we have taken into account Eq. (3). From Eq. (28), we obtain

$$- \int_{\mathcal{V}} d\mathbf{v} [\mathbf{v} - \mathbf{u}^\alpha(\mathbf{x})] \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) \ln \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}) = \frac{1}{T_\alpha} (\mathbf{J}_{\mathcal{E}_L} - \mathbf{J}_{\mathcal{E}_{Oh.}} - \mathbf{J}_{\mathcal{E}_{NBI}} - \mathbf{J}_{\mathcal{E}_{ICRH}}) \quad (30)$$

Here $\mathbf{J}_{\mathcal{E}_L}$ indicates the energy loss flux. $\mathbf{J}_{\mathcal{E}_{Oh.}}$, $\mathbf{J}_{\mathcal{E}_{NBI}}$ and $\mathbf{J}_{\mathcal{E}_{ICRH}}$ denote the Ohmic energy flux, the NBI energy flux and the ICRH energy flux, respectively. T_α is the temperature of species α . Eqs (24), (29) and (30), together with the definitions given by Eqs (21) and (22), allow determining the seven parameters $\Theta, \gamma, P_{\phi 0}, \lambda_0, \Delta P_\phi, \Delta \lambda_0, \Delta \lambda_1$. More particularly, the values of these parameters may be fixed by adopting the following strategy. The coefficients $\gamma, \Delta P_\phi, \Delta \lambda_0$ and $\Delta \lambda_1$ are provided by Eq. (29), whereas Eq. (24) sets up the calculations for the determination of parameter Θ . $P_{\phi 0}$ and λ_0 are determined by expansions (9). The plasma temperature profile can be obtained by Eq. (30). This solution should be in agreement with the definition given by Eq. (23). Generally, the latter calculation is very complex, but it can be strongly simplified by adopting the following procedure. In a first phase, the plasma is heated to thermonuclear conditions by the externally supplied power. In our simplified case, the applied sources are the sum of $\Delta \mathcal{E}_{Oh.}$, $\Delta \mathcal{E}_{NBI}$ and $\Delta \mathcal{E}_{ICRH}$. When the thermonuclear conditions have been achieved, the plasma temperature can be sustained if, in the overall power balance, the power loss, $\mathcal{P}_L$, is balanced by the external supplied Ohmic ($\mathcal{P}_{Oh.}$), NBI ($\mathcal{P}_{NBI}$), and ICRH ($\mathcal{P}_{ICRH}$) powers as well as the *alpha-power heating* ($\mathcal{P}_{alpha}$) i.e.,

$$\mathcal{P}_{alpha} + \mathcal{P}_{Oh.} + \mathcal{P}_{NBI} + \mathcal{P}_{ICRH} = \mathcal{P}_L \quad (31)$$

$$\mathbf{J}_S^\alpha(\mathbf{x}) = 0$$

Of course, Eq. (24) and Eq. (30) remain valid also for burning plasma, which can be used after having specified a proper model for estimating the source $\mathcal{P}_{alpha}$. However, as already mentioned, the loss due to the process of producing the radiation (Bremsstrahlung) is neglected and $\mathcal{P}_{alpha}$ is not treated in this work. Since the quantities entering in this equation depend on the plasma temperature, Eq. (31) may be solved for estimating the *average temperature*. However, as previously mentioned, calculations can be performed only after having modeled the external energy sources and estimated the entropy source strength

σ^α by modeling the magnetically confined plasmas. Below, we provide some examples of modeling.

The Ohmic Heating Model

The expression of the Ohmic heating density can be found in many reference books. For the magnetic configuration (49), we have (see, for example, Refs [16] and [18])

$$\mathcal{P}_{Oh.} = \eta j^2 = 2\eta \left[\frac{c^2 B_0}{4\pi R_0 (1 + r/R_0 \cos \theta)} \right]^2 \frac{1}{q(0)[q(a) - 1/2q(0)]} \quad (32)$$

$$\eta = \frac{\eta_S}{[(1 - r/R_0)^{1/2}]^2} \quad (33)$$

where η is the resistivity of the plasma and $\mathbf{j}$ is the current density. a and η_S denote the minor radius of the tokamak and the Spitzer resistivity, respectively.

The NBI Source Model

Once the neutral beam enters in the plasma, the neutral particles will be ionized. Their energy passes to the particles of the plasmas causing heating of both electrons and ions. The energy supplied by neutral beam injection may be modeled thinking in terms of a pencil beam source, S_{NBI} , acting for a period of time τ [19]

$$S_{NBI} = \dot{n}_b \tau \delta(\mathbf{v} - \mathbf{v}_b) \quad (34)$$

where $\dot{n}_{NBI}$ is the birth rate per unit volume and $\mathbf{v}_b$ is the beam ion velocity. Hence, the NBI energy is

$$\Delta \mathcal{E}_{NBI} \simeq \frac{1}{2} m_b \dot{n}_b \tau |\mathbf{v}_b|^2 \Omega_P \quad (35)$$

where m_b and Ω_P are the mass of the beam ion and the plasma volume, respectively.

The ICRH Source Model

Let us consider the case of a low concentration of ions ${}^3He$ colliding with a thermal background plasma, composed by Deuterium and electrons. The ${}^3He$ minority is about 2% – 3% of the density of the background plasma and it is heated by ion cyclotron resonant heating (ICRH). In the velocity space, the long term evolution of the distribution function for the high frequency heated ions, is governed by [20], (see the footnote [34])

$$\frac{\partial \mathcal{F}^m(\mathbf{y}, t)}{\partial t} = -\nabla \cdot \mathbf{S}^m(\mathcal{F}^m) + P(\mathcal{F}^m) \quad (36)$$

where

$$\mathbf{S}^m(\mathcal{F}^m) = \bar{\mathbf{S}}_W^m(\mathcal{F}^m) + \sum_{\alpha=e,i} \bar{\mathbf{S}}_c^{m\alpha}(\mathcal{F}^m) \quad (37)$$

Suffices m and α [with $m = {}^3\text{He}$ and $\alpha = (e, i)$] distinguish the minority population and the species of the background plasma, respectively. The first term in Eq. (37) describes the quasi-linear diffusion due to the resonant wave particle interactions and the second term is due to the collisional operator. $P(\mathcal{F}^m)$ takes into account other auxiliary sources; in our analysis, we shall put $P(\mathcal{F}^m) = 0$. The gradient operator, ∇ , is defined as the row vector $\nabla \equiv (\partial_{v_x}, \partial_{v_y}, \partial_{v_z})$ in the velocity space, whilst $\nabla \cdot \bar{\bar{\mathbf{A}}}$ is the matrix multiplication between the gradient vector and the matrix $\bar{\bar{\mathbf{A}}}$. In this approach, the ICRH is modeled by the divergence of the quasi-linear flux $\bar{\mathbf{S}}_W^m(\mathcal{F}^m)$, which is proportional to the energy density of the propagating wave

$$\Delta \mathcal{E}_{ICRH} \simeq -\frac{1}{2} m_m \tau \int_{\{\Omega \nu\}} d\Gamma |\mathbf{v} - \mathbf{u}_m|^2 \nabla \cdot \bar{\mathbf{S}}_W^m(\mathcal{F}^m) \quad (38)$$

where τ , m_m and $\mathbf{u}_m$ are the small laps of time during which the minority population is heated by ICRH, the mass and the average velocity of the minority, respectively. $\bar{\mathbf{S}}_c^{m\alpha}$ and the simplified quasi-linear term, $\bar{\mathbf{S}}_w^m$, can be written as [21], [22]

$$\begin{aligned} \bar{\mathbf{S}}_c^{m\alpha}(\mathcal{F}^m) &= -[\nabla \cdot (\bar{\bar{\mathbf{D}}}_c^{m\alpha(2)} \mathcal{F}^m)]^T + \bar{\mathbf{D}}_c^{m\alpha(1)} \mathcal{F}^m \\ \bar{\mathbf{S}}_W^m(\mathcal{F}^m) &= -[\nabla \cdot (\bar{\bar{\mathbf{D}}}_W^{m(2)} \mathcal{F}^m)]^T + \bar{\mathbf{D}}_W^{m(1)} \mathcal{F}^m \end{aligned} \quad (39)$$

with T denoting the *transpose operation*. We introduce the following dimensionless coordinates, $\hat{w}$ and $\hat{\lambda}$

$$\begin{aligned} \hat{w} &= \frac{w}{v_{the}^2} & \text{where } v_{th\alpha} &= \sqrt{\frac{2T_\alpha}{m_\alpha}} & \alpha &= (e, i) \\ \hat{\lambda} &= B_0 \lambda \end{aligned} \quad (40)$$

with w and λ defined in Eqs (16)]. In these coordinates, the matrices $\bar{\bar{\mathbf{D}}}_c^{m\alpha(2)}$, $\bar{\bar{\mathbf{D}}}_W^{m(2)}$, $\bar{\mathbf{D}}_c^{m\alpha(1)}$ and $\bar{\mathbf{D}}_W^{m(1)}$ can be cast into the form

$$\begin{aligned}\bar{\bar{\mathbf{D}}}_c^{me(2)} &= \begin{pmatrix} D_{cww}^{me(2)} & 0 \\ 0 & D_{c\lambda\lambda}^{me(2)} \end{pmatrix} \quad ; \quad \bar{\bar{\mathbf{D}}}_c^{mi(2)} = \begin{pmatrix} D_{cww}^{mi(2)} & 0 \\ 0 & D_{c\lambda\lambda}^{mi(2)} \end{pmatrix} \\ \bar{\mathbf{D}}_c^{me(1)} &= \begin{pmatrix} D_{cw}^{me(1)} + \frac{2}{v_{the}\sqrt{\hat{w}}} D_{cww}^{me(2)} + \frac{2\sqrt{\hat{w}}}{v_{the}} \frac{dD_{cww}^{me(2)}}{d\hat{w}} \\ \frac{1}{v_{the}} \sqrt{\frac{1-2|\hat{B}|\hat{\lambda}}{2|\hat{B}|\hat{\lambda}\hat{w}}} D_{c\lambda\lambda}^{me(2)} \end{pmatrix} \\ \bar{\mathbf{D}}_c^{mi(1)} &= \begin{pmatrix} D_{cw}^{mi(1)} + \frac{2}{v_{the}\sqrt{\hat{w}}} D_{cww}^{mi(2)} + \frac{2\sqrt{\hat{w}}}{v_{the}} \frac{dD_{cww}^{mi(2)}}{d\hat{w}} \\ \frac{1}{v_{the}} \sqrt{\frac{1-2|\hat{B}|\hat{\lambda}}{2|\hat{B}|\hat{\lambda}\hat{w}}} D_{c\lambda\lambda}^{mi(2)} \end{pmatrix}\end{aligned}\tag{41}$$

with $|\hat{B}| \equiv |B|/B_0$, and

$$\begin{aligned}\bar{\bar{\mathbf{D}}}_W^{m(2)} &= \begin{pmatrix} D_{Www}^{m(2)} & D_{Ww\lambda}^{m(2)} \\ D_{W\lambda w}^{m(2)} & D_{W\lambda\lambda}^{m(2)} \end{pmatrix} \\ \bar{\mathbf{D}}_W^{m(1)} &= \begin{pmatrix} \frac{2}{v_{the}\sqrt{\hat{w}}} D_{Www}^{m(2)} + \frac{2\sqrt{\hat{w}}}{v_{the}} \frac{dD_{Www}^{m(2)}}{d\hat{w}} + \frac{1}{v_{the}} \sqrt{\frac{1-2|\hat{B}|\hat{\lambda}}{2|\hat{B}|\hat{\lambda}\hat{w}}} D_{Ww\lambda}^{m(2)} + \frac{1}{v_{the}} \sqrt{\frac{2\hat{\lambda}(1-2|\hat{B}|\hat{\lambda})}{|\hat{B}|\hat{w}}} \frac{dD_{Ww\lambda}^{m(2)}}{d\hat{\lambda}} \\ \frac{2}{v_{the}\sqrt{\hat{w}}} D_{W\lambda w}^{m(2)} + \frac{2\sqrt{\hat{w}}}{v_{the}} \frac{dD_{W\lambda w}^{m(2)}}{d\hat{w}} + \frac{1}{v_{the}} \sqrt{\frac{1-2|\hat{B}|\hat{\lambda}}{2|\hat{B}|\hat{\lambda}\hat{w}}} D_{W\lambda\lambda}^{m(2)} + \frac{1}{v_{the}} \sqrt{\frac{2\hat{\lambda}(1-2|\hat{B}|\hat{\lambda})}{|\hat{B}|\hat{w}}} \frac{dD_{W\lambda\lambda}^{m(2)}}{d\hat{\lambda}} \end{pmatrix}\end{aligned}\tag{42}$$

Notice that in Eqs (41) and (42), extra terms appear in the components of the drift vectors. However, the nature of these additional terms is *purely geometrical* and they come from the conversion of the balance equation, Eq. (36), into the standard Fokker-Planck equation (see, for instance, Refs [23] or [24]). Hence, the expressions of the collisional electron and ion coefficients can be brought into the form

$$\begin{aligned}D_{cww}^{me(2)} &= \frac{\Gamma^{me}}{2v_{the}\sqrt{\hat{w}}} \left(\frac{\text{erf}(\sqrt{\hat{w}}) - \sqrt{\hat{w}} \text{erf}'(\sqrt{\hat{w}})}{\hat{w}} \right) \\ D_{c\lambda\lambda}^{me(2)} &= \frac{\Gamma^{me}}{4v_{the}\sqrt{\hat{w}}} \left(\left(2 - \frac{1}{\hat{w}} \right) \text{erf}(\sqrt{\hat{w}}) + \frac{\text{erf}'(\hat{w})}{\sqrt{\hat{w}}} \right) \\ D_{cw}^{me(1)} &= -\frac{\Gamma^{me}}{v_{the}^2} \frac{m_i}{m_e} \frac{[\text{erf}(\sqrt{\hat{w}}) - \sqrt{\hat{w}} \text{erf}'(\sqrt{\hat{w}})]}{\hat{w}} \quad \text{where} \\ \Gamma^{m\alpha} &= \frac{4\sqrt{2\pi}}{3} \frac{n_\alpha e^4 Z_m^2 Z_\alpha^2 \ln \Lambda_{m\alpha}}{m_m^2} \quad ; \quad \Lambda_{m\alpha} = \frac{3(T_m + T_\alpha) \lambda_{Dm\alpha}}{2Z_m Z_\alpha e^2} \\ \lambda_{Dm\alpha} &= \left(\frac{T_m T_\alpha (Z_m + Z_\alpha)}{4\pi Z_m Z_\alpha e^2 (n_m T_m + n_\alpha T_\alpha)} \right)^{1/2}\end{aligned}\tag{43}$$

and

$$\begin{aligned}
D_{cw}^{mi(2)} &= \frac{\Gamma^{mi}}{2v_{the}\sqrt{\hat{w}}} \left(\frac{\text{erf}(\sqrt{\kappa\hat{w}}) - \sqrt{\kappa\hat{w}} \text{erf}'(\sqrt{\kappa\hat{w}})}{\kappa\hat{w}} \right) \\
D_{c\lambda}^{mi(2)} &= \frac{\Gamma^{mi}}{4v_{the}\sqrt{\hat{w}}} \left(\left(2 - \frac{1}{\kappa\hat{w}}\right) \text{erf}(\sqrt{\kappa\hat{w}}) + \frac{\text{erf}'(\sqrt{\kappa\hat{w}})}{\sqrt{\kappa\hat{w}}} \right) \\
D_{cw}^{mi(1)} &= -\frac{\Gamma^{mi}}{v_{the}^2} \frac{(\text{erf}(\sqrt{\kappa\hat{w}}) - \sqrt{\kappa\hat{w}} \text{erf}'(\sqrt{\kappa\hat{w}}))}{\hat{w}} \quad ; \quad \text{where} \quad \kappa \equiv \frac{v_{the}^2}{v_{thi}^2}
\end{aligned} \tag{44}$$

respectively. Notice that, the quantities in Eqs (43), are expressed in the *CGS Gaussian units*. $\text{erf}(x)$ and $\text{erf}'(x)$ denote the *error function* and the *derivative* of the error function with respect to its argument, respectively. The expressions of the diffusion coefficients for the resonant wave particle interactions are [21], [22]

$$\begin{aligned}
D_{Ww}^{m(2)} &= 2 \mid \hat{B} \mid \hat{\lambda} D_{\perp\perp}^m \\
D_{Ww\lambda}^{m(2)} &= D_{W\lambda w}^{m(2)} = \sqrt{(2 \mid \hat{B} \mid \hat{\lambda})(1 - 2 \mid \hat{B} \mid \hat{\lambda})} D_{\perp\perp}^m \\
D_{W\lambda\lambda}^{m(2)} &= (1 - 2 \mid \hat{B} \mid \hat{\lambda}) D_{\perp\perp}^m \\
D_{\perp\perp}^m &= D_0^m J_p^2(\xi_m \sqrt{2 \mid \hat{B} \mid \hat{\lambda} \hat{w}}) \quad \text{with} \\
\xi_m &= \frac{v_{the}\kappa_{\perp}}{\Omega_{cm}} \quad ; \quad \Omega_{cm} = \frac{Z_m e B_0}{m_m c} \quad ; \quad P_{abs}^{lin} \simeq 10 \div 80 \text{ W/cm}^3 \\
D_0^m &= \frac{P_{abs}^{lin}}{4m_m n_m \int_0^\infty x^3 J_p^2(\xi_m x) \exp(-x^2) dx}
\end{aligned} \tag{45}$$

Here, $J_p(x)$ indicates the Bessel functions of the first kind. e and c are the absolute value of the charge of the electron and the speed of light, respectively. From Eqs (41)-(45) we obtain the expressions of the *total matrix diffusion coefficients* and the *total drift vector coefficients*

$$\begin{aligned}
\bar{\mathbf{D}}^{m(2)} &= \bar{\mathbf{D}}_c^{me(2)} + \bar{\mathbf{D}}_c^{mi(2)} + \bar{\mathbf{D}}_W^{m(2)} = \begin{pmatrix} D_{ww}^{m(2)} & D_{w\lambda}^{m(2)} \\ D_{w\lambda}^{m(2)} & D_{\lambda\lambda}^{m(2)} \end{pmatrix} \\
\bar{\mathbf{D}}^{m(1)} &= \bar{\mathbf{D}}_c^{me(1)} + \bar{\mathbf{D}}_c^{mi(1)} + \bar{\mathbf{D}}_W^{m(1)} = \begin{pmatrix} D_w^{m(1)} \\ D_\lambda^{m(1)} \end{pmatrix}
\end{aligned} \tag{46}$$

In the zero orbit width limit, the steady state DDF of the Fokker-Planck equation, with the diffusion and drift coefficients given by Eqs (46), may be fitted by the profile

$$\mathcal{F}^m \propto \left(\frac{w}{\Theta}\right)^{-3/4} \exp\left[-\left(\frac{P - P_\phi}{\Delta P_\phi}\right)^2\right] \exp\left(-\frac{w}{\Theta}\right) \delta(\lambda - \lambda_0) \mid \mathcal{J} \mid \tag{47}$$




FIG. 7: Dimensionless diffusion component $\hat{D}_{ww}^{m(2)}$ versus the normalized, dimensionless variables $\hat{w}$ and $\hat{\lambda}$. The dimensionless diffusion coefficients are defined as $\hat{D}_{ij}^{m(2)} \equiv D_{ij}^{m(2)} v_{the} / \Gamma^{me}$.




FIG. 8: Dimensionless diffusion component $\hat{D}_{w\lambda}^{m(2)}$ versus $\hat{w}$ and $\hat{\lambda}$. The matrix of the diffusion coefficients is symmetric: $\hat{D}_{w\lambda}^{m(2)} = \hat{D}_{\lambda w}^{m(2)}$.

with $\delta(\lambda - \lambda_0)$ indicating the Dirac function [25]. Figs (7), (8) and (9), report on the graphics of the dimensionless elements of the total diffusion matrix, $\hat{D}_{ww}^{m(2)}$, $\hat{D}_{w\lambda}^{m(2)} = \hat{D}_{\lambda w}^{m(2)}$ and $\hat{D}_{\lambda\lambda}^{m(2)}$ [see the first equation in Eqs (46)], versus $\hat{w}$ and $\hat{\lambda}$, respectively. The dimensionless diffusion coefficients are defined as $\hat{D}_{ij}^{m(2)} \equiv D_{ij}^{m(2)} v_{the} / \Gamma^{me}$. Figs (10)-(11) report on the dimensionless components of the total drift coefficients [see the second equation in Eqs (46)],




FIG. 9: Dimensionless diffusion component $\hat{D}_{\lambda\lambda}^{m(2)}$ versus $\hat{w}$ and $\hat{\lambda}$.




FIG. 10: Plot of the first dimensionless component of the total drift vector, $\hat{D}_w^{m(1)}$, versus $\hat{w}$ and $\hat{\lambda}$.

defined as $\hat{D}_i^{m(1)} \equiv D_i^{m(1)} v_{the}^2 / \Gamma^{me}$, against $\hat{w}$ and $\hat{\lambda}$. These pictures have been obtained by considering only the fundamental cyclotron heating, $p = 0$, (minority heating) and by setting $\kappa_{\perp} \simeq \rho_{Lm}^{-1}$ (with ρ_{Lm} denoting the *Larmor gyroradius*). Moreover, as an example




FIG. 11: Plot of the second dimensionless component of the total drift vector, $\hat{D}_\lambda^{m(1)}$, versus $\hat{w}$ and λ .

of calculation, we chose the following values of the tokamak parameters: $B_0 = 3.45 \text{ T}$, $R_0 = 2.96 \text{ m}$ and $a = 1.25 \text{ m}$. Moreover, $P_{abs}^{lin} = 50 \text{ W/cm}^3$ and $n_m = 2.5\% n_i$, respectively and, as an example of calculation, the fields have been estimated at the center of the tokamak. Notice that in our case, the charge number and the ion mass are $Z_\alpha = 1$ and $m_i = m_D = 2m_H$ for the background, and $Z_m = Z_{^3\text{He}} = 2$ and $m_m = m_{^3\text{He}} \simeq 3m_H$ (with m_H indicating the mass of proton) for the minority, respectively.

ThePower Loss

Neglecting the Bremsstrahlung loss, the rate of energy loss is mainly due to the thermal conduction losses. Its expression can be found in many reference books [see, for example, Refs [16] and [18]]. We have

$$\mathcal{P}_L = 3\Omega_P \frac{\bar{n}\bar{T}}{\tau_E} \quad (48)$$

where $\bar{n}$ and $\bar{T}$ are the average density and the average temperature, respectively. τ_E is the *energy confinement time*.

Modeling Tokamak-Plasmas

The entropy source strength σ^α can be estimated by modeling the tokamak-plasma. For instance, let us consider fully ionized tokamak-plasmas, defined as a collection of magnetically confined electrons and positively charged ions. In the *local triad* ($\mathbf{e}_r, \mathbf{e}_\theta, \mathbf{e}_\phi$), the magnetic field, in the standard high aspect ratio, low *beta* (the plasma pressure normalized to the magnetic field strength), circular tokamak equilibrium model, reads (see, for example, Ref. [12]) (see the footnote [35])

$$\mathbf{B} = \frac{B_0}{q(r)} \frac{r}{R_0} \mathbf{e}_\theta + \frac{B_0}{1 + (r/R_0)\cos\theta} \mathbf{e}_\phi \quad (49)$$

Here B_0 is a constant having the dimension of a magnetic field intensity, and $q(r)$ and R_0 are the safety factor (in Ref. [12] the reader can find an exact definition of this tokamak parameter) and the major radius of the tokamak, respectively. In the magnetic configuration, given by Eq. (49), we have

$$F = B_0 R_0 \quad ; \quad \psi(r) = 2\pi B_0 \int_0^r \frac{r}{q(r)} dr \quad (50)$$

The dimensionless entropy production of species α , $\Delta_I S^\alpha$, is derived under the sole assumption that the state of the quiescent plasma is not too far from the reference local Maxwellian. In the *local dynamical triad* [26], $\Delta_I S^\alpha$ can be brought into the form (see Refs [12] and [27])

$$\begin{aligned} \Delta_I S^e &= q_{\parallel ps}^{(1)}(g_{\parallel}^{(1)} - \bar{g}_{\parallel}^{e(1)}) + q_{\parallel ps}^{e(3)}(g_{\parallel}^{e(3)} + \bar{g}_{\parallel}^{e(3)}) + q_{\parallel b}^{(1)}(g_{\parallel}^{(1)} - \bar{g}_{\parallel}^{e(1)}) + q_{\parallel b}^{e(3)}(g_{\parallel}^{e(3)} + \bar{g}_{\parallel}^{e(3)}) \\ &\quad + q_{\parallel b}^{e(5)} \bar{g}_{\parallel}^{e(5)} + \hat{q}_{\rho cl}^{e(1)} g_{\rho}^{(1)P} + \hat{q}_{\rho cl}^{e(3)} g_{\rho}^{e(3)} \\ \Delta_I S^i &= q_{\parallel ps}^{i(3)}(g_{\parallel}^{i(3)} + \bar{g}_{\parallel}^{i(3)}) + q_{\parallel b}^{i(3)}(g_{\parallel}^{i(3)} + \bar{g}_{\parallel}^{i(3)}) + q_{\parallel b}^{i(5)} \bar{g}_{\parallel}^{i(5)} + \hat{q}_{\rho cl}^{i(3)} g_{\rho}^{i(3)} \end{aligned} \quad (51)$$

Here $q_r^{\alpha(n)}$ [with $r = (\rho, \parallel, \wedge)$] denote the *Hermitian moments of the distribution functions* and $g_r^{\alpha(n)}$, $\bar{g}_r^{\alpha(n)}$ are the *dimensionless source terms*. Index n takes the values $n = (1, 3, 5)$. In the linear Onsager region, and up to the second order of the drift parameter ϵ , it can be shown that Eqs (51) simplify to [12]

$$\begin{aligned} \Delta_I S^e &= \tilde{\sigma}_{\parallel}(g_{\parallel}^{(1)} - \bar{g}_{\parallel}^{e(1)})^2 + \tilde{\kappa}_{\parallel}^e(g_{\parallel}^{e(3)} + \bar{g}_{\parallel}^{e(3)})^2 + \tilde{\epsilon}_{\parallel}^e(\bar{g}_{\parallel}^{e(5)})^2 + 2\tilde{\alpha}_{\parallel}(g_{\parallel}^{(1)} - \bar{g}_{\parallel}^{e(1)})(g_{\parallel}^{e(3)} + \bar{g}_{\parallel}^{e(3)}) \\ &\quad + 2\tilde{\gamma}_{\parallel}(g_{\parallel}^{(1)} - \bar{g}_{\parallel}^{e(1)})\bar{g}_{\parallel}^{e(5)} + 2\tilde{\delta}_{\parallel}^e(g_{\parallel}^{e(3)} + \bar{g}_{\parallel}^{e(3)})\bar{g}_{\parallel}^{e(5)} \\ &\quad + \tilde{\sigma}_{\perp}(g_{\rho}^{(1)P})^2 + \tilde{\kappa}_{\perp}^e(g_{\rho}^{e(3)})^2 - 2\tilde{\alpha}_{\perp} g_{\rho}^{(1)P} g_{\rho}^{e(3)} \\ \Delta_I S^i &= \tilde{\kappa}_{\parallel}^i(g_{\parallel}^{i(3)} + \bar{g}_{\parallel}^{i(3)})^2 + \tilde{\epsilon}_{\parallel}^i(\bar{g}_{\parallel}^{i(5)})^2 + 2\tilde{\delta}_{\parallel}^i(g_{\parallel}^{i(3)} + \bar{g}_{\parallel}^{i(3)})\bar{g}_{\parallel}^{i(5)} + \tilde{\kappa}_{\perp}^i(g_{\rho}^{i(3)})^2 \end{aligned} \quad (52)$$

where coefficients $\tilde{\sigma}_r$, $\tilde{\alpha}_r$, $\tilde{\kappa}_r^\alpha$ indicate the dimensionless component of the *electronic conductivity*, the *thermoelectric coefficient* and the *electric* ($\alpha = e$) or *ion* ($\alpha = i$) *thermal*

conductivity, respectively. Moreover, $\tilde{\gamma}_{\parallel}$, $\tilde{\delta}_{\parallel}^{\alpha}$ and $\tilde{\epsilon}_{\parallel}^{\alpha}$ are the *parallel transport coefficients* in 21M approximation. As shown in Eq. (29), the entropy production is closely associated with the collision term. Notice that Eqs. (52) have been derived by using the following *linearized Landau collisional term*

$$\begin{aligned} \mathcal{K}_L^{\alpha\beta} = & \frac{2\pi e_{\alpha}^2 e_{\beta}^2 \ln \Lambda}{m_{\alpha}} \int d\mathbf{v}_2 \frac{\partial}{\partial v_{1r}} G_{rs}(\mathbf{v}_1 - \mathbf{v}_2) \left(f_0^{\beta}(\mathbf{v}_2) \frac{1}{m_{\alpha}} \frac{\partial}{\partial v_{1s}} f_1^{\alpha}(\mathbf{v}_1) \right. \\ & \left. + f_1^{\beta}(\mathbf{v}_2) \frac{1}{m_{\alpha}} \frac{\partial}{\partial v_{1s}} f_0^{\alpha}(\mathbf{v}_1) - f_0^{\alpha}(\mathbf{v}_1) \frac{1}{m_{\beta}} \frac{\partial}{\partial v_{2s}} f_0^{\beta}(\mathbf{v}_2) - f_1^{\alpha}(\mathbf{v}_1) \frac{1}{m_{\beta}} \frac{\partial}{\partial v_{2s}} f_0^{\beta}(\mathbf{v}_2) \right) \end{aligned} \quad (53)$$

Moreover, $G_{rs}(\mathbf{g})$ and $\ln \Lambda$ denote the *Landau tensor* and the *Coulomb logarithm*, respectively

$$G_{rs}(\mathbf{g}) = \frac{g^2 \delta_{rs} - g_r g_s}{g^3} \quad ; \quad \ln \Lambda = \frac{3/2(T_e + T_i)\lambda_D}{Ze^2} \quad ; \quad \lambda_D = \left(\frac{4\pi Ze^2(n_e T_e + n_i T_i)}{T_e T_i (1 + Z)} \right)^{-1/2} \quad (54)$$

Here Z is the *charge number of ions*. In addition, in Eq. (53), the distribution function is expanded in powers of the drift parameter ϵ

$$\begin{aligned} f^{\alpha}(\mathbf{v}) &= f_0^{\alpha}(\mathbf{v}) + \epsilon f_1^{\alpha}(\mathbf{v}) + \dots \quad \text{with} \\ f_0^{\alpha}(\mathbf{v}) &= n_{\alpha} \left(\frac{m_{\alpha}}{2\pi T_{\alpha}} \right)^{3/2} \exp\left(-\frac{m_{\alpha}}{2T_{\alpha}} v^2\right) \end{aligned} \quad (55)$$

By summarizing, the expression of the entropy source strength, Eq. (52), may be put on the left-hand-side of Eq. (29) provide that the integral appearing on the right-hand-side of this equation is evaluated by means of the linearized Landau operator Eq. (53).

IV. ESTIMATION OF THE SEVEN PARAMETERS APPEARING IN THE EXPRESSION OF $\mathcal{F}^0$ FOR THE SIMPLE MODEL OF FULLY IONIZED TOKAMAK-PLASMAS

As an example of calculation we estimate now the seven parameters of $\mathcal{F}^0$ when the entropy production is given by Eq. (52). In particular, our purpose is to link the expressions of the coefficients $g_{j\kappa}$, appearing in Eq. (7), in terms of the transport coefficients. In addition we shall prove that the expression that we found for the reference DDF, $\mathcal{F}^0$, *coincides exactly* with the one predicted by the neoclassical theory for collisional tokamak-plasmas.

From Eqs (2), (7) and (13), we have

$$\Delta_I S = -(\gamma - 1) \ln\left(\frac{w}{\Theta}\right) + \frac{w}{\Theta} + \frac{1}{2} g_{11} \alpha_1^2 + \frac{1}{2} g_{22} \alpha_2^2 + g_{12} \alpha_1 \alpha_2 \quad (56)$$

By expanding the previous expression around the reference value $w = w_0$ we obtain, up to the second order

$$\begin{aligned}\Delta_I S = & \left[-(\gamma - 1) \ln\left(\frac{w_0}{\Theta}\right) + \frac{w_0}{\Theta} \right] + \left[-(\gamma - 1) \frac{1}{w_0} + \frac{1}{\Theta} \right] (w - w_0) + (\gamma - 1) \frac{1}{2w_0^2} (w - w_0)^2 \\ & + \frac{1}{2} g_{11} \alpha_1^2 + \frac{1}{2} g_{22} \alpha_2^2 + g_{12} \alpha_1 \alpha_2 + h.o.t.\end{aligned}\quad (57)$$

Up to a normalization constant, we have that the distribution function Eq. (19) is approximated by a Gaussian density distribution function (in the variable w) by setting to zero the coefficient of the linear term [i.e., $-(\gamma - 1)/w_0 + 1/\Theta = 0$]. The *global optimality conditions* is obtained by imposing that also the sum of the constant terms vanishes [i.e., $-(\gamma - 1) \ln(w_0/\Theta) + w_0/\Theta = 0$]. These two requirements are simultaneously satisfied only if

$$\gamma = 1 + E \quad (58)$$

$$w_0 = (\gamma - 1)\Theta = E\Theta$$

Hence, solutions (58) ensure not only a local approximation, valid up to the second order, but also a good global approximation. In this sense the values of w_0 and γ , provided by Eq. (58), are *optimal*. From Eq. (4) and Eq. (57) we obtain the expressions of the thermodynamic forces

$$\begin{aligned}X^1 &\equiv \frac{\partial \Delta_I S}{\partial \alpha_1} = g_{11} \alpha_1 + g_{12} \alpha_2 \\ X^2 &\equiv \frac{\partial \Delta_I S}{\partial \alpha_2} = g_{12} \alpha_1 + g_{22} \alpha_2 \\ X^3 &\equiv \frac{\partial \Delta_I S}{\partial w} = \frac{1}{E\Theta^2} (w - w_0)\end{aligned}\quad (59)$$

By solving the previous system of equations with respect to α_1 , α_2 and $(w - w_0)$, we find

$$\begin{aligned}\alpha_1 &= \hat{g}_{22} X^1 - \hat{g}_{12} X^2 \\ \alpha_2 &= -\hat{g}_{12} X^1 + \hat{g}_{11} X^2 \\ (w - w_0) &= E\Theta^2 X^3 \quad \text{where} \\ \hat{g}_{j\kappa} &\equiv \frac{g_{j\kappa}}{g} \quad [\text{with } (j, \kappa) = (1, 2)] \quad ; \quad g \equiv g_{11}g_{22} - g_{12}^2 = \frac{1}{\hat{g}_{11}\hat{g}_{22} - \hat{g}_{12}^2} = \frac{1}{\hat{g}}\end{aligned}\quad (60)$$

Hence, in terms of the thermodynamic forces, the electron and ion entropy source strength read, respectively

$$\begin{aligned}\Delta_I S^e &= \frac{1}{2} E \Theta_e^2 X_e^{3^2} + \frac{1}{2} \hat{g}_{22}^e X_e^{1^2} + \frac{1}{2} \hat{g}_{11}^e X_e^{2^2} - \hat{g}_{12}^e X_e^1 X_e^2 + h.o.t. \\ \Delta_I S^i &= \frac{1}{2} E \Theta_i^2 X_i^{2^2} + \frac{1}{2} \hat{g}_{11}^i X_i^{1^2} + h.o.t.\end{aligned}\quad (61)$$

By diagonalizing, the first expression of Eq. (61) can be brought into the form

$$\begin{aligned}
\Delta_I S^e &= \frac{1}{2} E \Theta_e^2 X_e^{3^2} + \mu_1 \xi_e^{1^2} + \mu_2 \xi_e^{2^2} \quad \text{with} \\
\mu_{1,2} &= \frac{1}{4} \left[(\hat{g}_{11} + \hat{g}_{22}) \pm \sqrt{(\hat{g}_{11} - \hat{g}_{22})^2 + 4\hat{g}_{12}^2} \right] \\
\xi_e^1 &= \frac{q_1}{h} \hat{g}_{12} X_e^1 - \frac{q_1}{2h} (\hat{g}_{11} - \hat{g}_{22} - h) X_e^2 \\
\xi_e^2 &= -\frac{q_2}{h} \hat{g}_{12} X_e^1 + \frac{q_2}{2h} (\hat{g}_{11} - \hat{g}_{22} + h) X_e^2 \\
h &= \sqrt{(\hat{g}_{11} - \hat{g}_{22})^2 + 4\hat{g}_{12}^2} \\
q_{1,2} &= \frac{\sqrt{4\hat{g}_{12}^2 + (\hat{g}_{11} - \hat{g}_{22})(\hat{g}_{11} - \hat{g}_{22} \pm h)}}{|\hat{g}_{12}| \sqrt{2}}
\end{aligned} \tag{62}$$

Let us now evaluate the coefficients $\hat{g}_{j\kappa}$ and Θ_α for the case of fully ionized tokamak-plasmas described, for example, in Ref. [12]. In the banana regime, without considering the classical contributions, the expressions of the electron and ion entropy source strength take the form

$$\begin{aligned}
\Delta_I S^{e'} &= \tilde{\epsilon}_\parallel^e Z_e^{1^2} + \tilde{\kappa}_\parallel^e Z_e^{2^2} + \tilde{\sigma}_\parallel Z_e^{3^2} + 2\tilde{\delta}_\parallel^e Z_e^1 Z_e^2 + 2\tilde{\gamma}_\parallel Z_e^1 Z_e^3 + 2\tilde{\alpha}_\parallel Z_e^2 Z_e^3 \\
\Delta_I S^{i'} &= \tilde{\epsilon}_\parallel^i Z_i^{1^2} + \tilde{\kappa}_\parallel^i Z_i^{2^2} + 2\tilde{\delta}_\parallel^i Z_i^1 z_i^2
\end{aligned} \tag{64}$$

where

$$\left\{ \begin{array}{l} Z_e^1 = \bar{g}_\parallel^{e(5)} \\ Z_e^2 = g_\parallel^{e(3)} + \bar{g}_\parallel^{e(3)} \\ Z_e^3 \equiv g_\parallel^{(1)} - \bar{g}_\parallel^{e(1)} \end{array} \right. ; \quad \left\{ \begin{array}{l} Z_i^1 = g_\parallel^{i(3)} + \bar{g}_\parallel^{i(3)} \\ Z_i^2 = \bar{g}_\parallel^{i(5)} \end{array} \right. \tag{65}$$

As mentioned in the introduction [see Eq. (8)], there exists the following (linear) thermodynamic covariant transformation (TCT) (see Eq. (6) and Ref. [8])

$$\left\{ \begin{array}{l} X_e^1 = Z_e^1 = \bar{g}_\parallel^{e(5)} \\ X_e^2 = Z_e^2 = g_\parallel^{e(3)} + \bar{g}_\parallel^{e(3)} \\ X_e^3 \equiv a_e Z_e^1 + b_e Z_e^2 + c_e Z_e^3 \end{array} \right. ; \quad \left\{ \begin{array}{l} X_i^1 = Z_i^1 = g_\parallel^{i(3)} + \bar{g}_\parallel^{i(3)} \\ X_i^2 = a_i Z_i^1 + b_i Z_i^2 \end{array} \right. \tag{66}$$

such that

$$\begin{aligned}
\Delta_I S^e &= \Delta_I S^{e'} \\
\Delta_I S^i &= \Delta_I S^{i'}
\end{aligned} \tag{67}$$

The transport coefficients $\hat{g}_{ij}^\alpha$ and the transformation-coefficients a_α, b_α [with $\alpha = (e, i)$] and

c_e , are easily obtained from Eqs (61), (64) and Eqs (67). We find

$$\begin{aligned}
a_e &= \frac{2\tilde{\gamma}_{\parallel}}{\sqrt{E\tilde{\sigma}_{\parallel}}\Theta_e} \quad ; \quad b_e = \frac{2\tilde{\alpha}_{\parallel}}{\sqrt{E\tilde{\sigma}_{\parallel}}\Theta_e} \quad ; \quad c_e = \frac{2\sqrt{\tilde{\sigma}_{\parallel}}}{\sqrt{E}\Theta_e} \\
\hat{g}_{11}^e &= \frac{2}{\tilde{\sigma}_{\parallel}}(\tilde{\kappa}_{\parallel}^e\tilde{\sigma}_{\parallel} - \tilde{\alpha}_{\parallel}^2) \quad ; \quad \hat{g}_{22}^e = \frac{2}{\tilde{\sigma}_{\parallel}}(\tilde{\epsilon}_{\parallel}^e\tilde{\sigma}_{\parallel} - \tilde{\gamma}_{\parallel}^2) \quad ; \quad \hat{g}_{12}^e = \frac{2}{\tilde{\sigma}_{\parallel}}(\tilde{\alpha}_{\parallel}\tilde{\gamma}_{\parallel} - \tilde{\delta}_{\parallel}^e\tilde{\sigma}_{\parallel}) \\
a_i &= \frac{2\tilde{\delta}_{\parallel}^i}{\sqrt{E\tilde{\kappa}_{\parallel}^i}\Theta_i} \quad ; \quad b_i = \frac{2\sqrt{\tilde{\kappa}_{\parallel}^i}}{\sqrt{E}\Theta_i} \quad ; \quad \hat{g}_{11}^i = \frac{2}{\tilde{\kappa}_{\parallel}^i}(\tilde{\epsilon}_{\parallel}^i\tilde{\kappa}_{\parallel}^i - \tilde{\delta}_{\parallel}^{i2})
\end{aligned} \tag{68}$$

By inserting the values of the collision matrix elements (see, for example, Ref. [6]), we find

$$\begin{aligned}
a_e &= \frac{0.1168}{\Theta_e} \quad ; \quad b_e = -\frac{0.7599}{\Theta_e} \quad ; \quad c_e = \frac{1.6943}{\Theta_e} \\
\hat{g}_{11}^e &= 2.5125 \quad ; \quad \hat{g}_{22}^e = 1.9090 \quad ; \quad \hat{g}_{12}^e = -1.6583 \\
a_i &= \frac{0.6364}{\Theta_i} \quad ; \quad b_i = \frac{1.8033}{\Theta_i} \quad ; \quad \hat{g}_{11}^i = 2.1994
\end{aligned} \tag{69}$$

The numerical values of parameters Θ_{α} can be obtained by means of Eqs (21), (22) and (24) where the distribution functions $\mathcal{F}^{\alpha 0}$ are constructed by using Eqs (61). From Eq. (21) we obtain an equation, depending on variable Θ_{α} , for the particle density n_{α} .

$$n_{\alpha}(\mathbf{x}, \Theta_{\alpha}) = \int d\mathbf{v} \mathcal{F}^{\alpha 0}(\Theta_{\alpha}, \mathbf{v}, \mathbf{x}) \quad ; \quad \alpha = (e, i) \tag{70}$$

By injecting Eq. (70) into Eq. (22), we have

$$\mathbf{u}_{\alpha}(\mathbf{x}, \Theta_{\alpha}) = \frac{\int d\mathbf{v} \mathbf{v} \mathcal{F}^{\alpha 0}(\Theta_{\alpha}, \mathbf{v}, \mathbf{x})}{\int d\mathbf{v} \mathcal{F}^{\alpha 0}(\Theta_{\alpha}, \mathbf{v}, \mathbf{x})} \tag{71}$$

This latter equation should then be combined with Eq. (24) and, for a given value of $\Delta\mathcal{E}_{Oh.}$, we finally derive the equation for θ_{α}

$$\frac{1}{2}m_{\alpha} \int_{\Omega} d\mathbf{\Gamma} \mid \mathbf{v} - \mathbf{u}_{\alpha}(\mathbf{x}, \Theta_{\alpha}) \mid^2 \mathcal{F}^{\alpha 0}(\mathbf{v}, \mathbf{x}, \Theta_{\alpha}) = \Delta\mathcal{E}_{Oh.} \quad ; \quad \alpha = (e, i) \tag{72}$$

with $\mathbf{u}_{\alpha}$ is provided by Eq. (71). Eq. (72) can be solved (numerically) with respect to variable Θ_{α} ; we find

$$(\Theta_e, \Theta_i) = (4.1760 \times 10^{18} cm^2 sec^{-2}, 2.2745 \times 10^{15} cm^2 sec^{-2}) \tag{73}$$

From Eq. (70), Eq. (71) and Eq. (23), we can now derive the profiles of the particle density, $n_{\alpha}(\mathbf{x})$, the average velocity, $\mathbf{u}_{\alpha}$, and temperature, T_{α} , respectively. The knowledge of the fields n_{α} , T_{α} etc. allows determining the profiles of the thermodynamic forces Z_{α}^j [j=(1,2,3)]

for electrons, and $j=(1,2)$ for ions]. Parameters $P_{\phi 0}$ and λ_0 are determined by Eqs (9). By substituting Eqs (16) into Eq. (19), we can easily check that distribution function, Eq. (19), tends to reduce to a pure gamma-process as $P_\phi \rightarrow \psi(r)$, $\lambda \rightarrow (2|B|)^{-1}$ and $v_\parallel \rightarrow 0$ (i.e., the pitch angle θ_P is close to $\pi/2$). This happens when the thermodynamic forces, X^1 and X^2 , tend to vanish. Our aim is to derive the expressions of the steady-state electron and ion distribution functions for plasmas in the Onsager region, and $v_\parallel$, w , P_ϕ and λ close to 0, $E\Theta$, $\psi(r)$ and $[2|B|(r, \theta)]^{-1}$ respectively (see also the footnote [36]). To this purpose, we shall work with the dimensionless variables $\hat{P}_\phi$ and $\hat{\lambda}$, which for the magnetic configuration given by Eq. (49), are defined as

$$\begin{aligned}\hat{P}_\phi &\equiv \frac{1}{B_0 a^2} P_\phi = \frac{B_0 R_0}{a^2 \Omega_{0c}} \frac{v_\parallel}{|B|} + 2\pi \int_0^\rho \frac{\rho'}{q(\rho')} d\rho' \\ \hat{\lambda} &\equiv B_0 \lambda = \frac{B_0}{2|B|} \left(1 - \frac{v_\parallel^2}{w}\right) \\ \text{with } \rho &\equiv \frac{r}{a} \quad ; \quad |B| \simeq B_0 \left(1 - \frac{a}{R_0} \rho \cos \theta\right)\end{aligned}\tag{74}$$

We have

$$\begin{aligned}\hat{P}_{\phi 0}^\alpha &= \hat{P}_\phi^\alpha |_{v_\parallel \rightarrow 0} = 2\pi \int_0^{\rho_{\alpha 0}} \frac{\rho'}{q(\rho')} d\rho' = \hat{\psi}(\rho_{\alpha 0}) \quad ; \quad \alpha = (e, i) \\ \hat{\lambda}_0^\alpha &= \hat{\lambda}^\alpha |_{v_\parallel \rightarrow 0} = \frac{B_0}{2|B|} \simeq \frac{1}{2} \left(1 + \frac{a}{R_0} \rho_{\alpha 0} \cos \theta_{\alpha 0}\right) \quad ; \quad \alpha = (e, i)\end{aligned}\tag{75}$$

where (ρ_{e0}, θ_{e0}) and (ρ_{i0}, θ_{i0}) are solutions of the equations (see the footnote [37])

$$\begin{cases} X_e^1(r, \theta) = g_\parallel^{e(3)} + \bar{g}_\parallel^{e(3)} = \mathcal{O}(\epsilon) \\ X_e^2(r, \theta) = \bar{g}_\parallel^{e(5)} = \mathcal{O}(\epsilon) \end{cases} \quad ; \quad \begin{cases} X_i^1(r, \theta) = g_\parallel^{i(3)} + \bar{g}_\parallel^{i(3)} = \mathcal{O}(\epsilon) \\ r_{i0} \in \{r \mid x_i^{(1)}(r, \theta) = \mathcal{O}(\epsilon)\} \end{cases}\tag{76}$$

with the drift parameter ϵ of the order $\epsilon \simeq 10^{-3}$ [12]. As an example of calculation, we choose the following values of the parameters: $B_0 = 3.45 \text{ Tesla}$, $R_0 = 2,96 \text{ m}$ and $a = 1,25 \text{ m}$. In this case, the systems (76) admits the solutions

$$\begin{cases} (\rho_{e0}, \theta_{e0}) = (0.2376, 1.0562) \\ X_e^1 = X_e^2 = 2.0367 \times 10^{-4} \sim \mathcal{O}(\epsilon) \end{cases} \quad ; \quad \begin{cases} (\rho_{i0}, \theta_{i0}) = (0.5079, 2.8137) \\ X_i^1 = 0 \end{cases}\tag{77}$$

Fig. (12) illustrates the surfaces of the electron forces X_e^1 and X_e^2 . The intersection line corresponds to the values of (ρ, θ) such that $X_e^1 = X_e^2$. Fig. (13) shows the curve $\rho = \rho(\theta)$ where the ion thermodynamic force X_i^1 vanishes. The values of parameters $P_{\phi 0}^\alpha$ and λ_0^α are easily estimated from Eqs (75)




FIG. 12: The electron thermodynamic forces X_e^1 and X_e^2 . The intersection line corresponds to the values (ρ, θ) for which $X_e^1 = X_e^2$. The first system of equations in Eqs (76), is satisfied in the narrow region around to the values $(\rho_{e0}, \theta_{e0}) = (0.2376, 1.0562)$.




FIG. 13: Locus of the points (r, θ) where $X_i^1 = 0$. As an example of calculation, we choose the point $(\rho_{i0}, \theta_{i0}) = (0.5079, 2.8137)$ of this curve.

$$(\hat{P}_{\phi 0}^e, \hat{\lambda}_0^e) = (0.1651, 0.5246) \quad ; \quad (\hat{P}_{\phi 0}^i, \hat{\lambda}_0^i) = (0.5499, 0.3984) \quad (78)$$

The parameters $\Delta P_\phi, \Delta \lambda_0, \Delta \lambda_1$ can be obtained from the relations

$$\begin{aligned} \left(\frac{\partial X_e^1}{\partial \rho} \bigg|_{\rho_{e0}, \theta_{e0}}, \frac{\partial X_e^1}{\partial \theta} \bigg|_{\rho_{e0}, \theta_{e0}} \right) &= (0.0039, 1.5957 \times 10^{-4}) \\ \left(\frac{\partial X_e^2}{\partial \rho} \bigg|_{\rho_{e0}, \theta_{e0}}, \frac{\partial X_e^2}{\partial \theta} \bigg|_{\rho_{e0}, \theta_{e0}} \right) &= (-0.0028, -7.3501 \times 10^{-5}) \\ \left(\frac{\partial X_i^1}{\partial \rho} \bigg|_{\rho_{i0}, \theta_{i0}}, \frac{\partial X_i^1}{\partial \theta} \bigg|_{\rho_{i0}, \theta_{i0}} \right) &= (6.6596 \times 10^{-4}, \mathcal{O}(\epsilon^2)) \end{aligned} \quad (79)$$

By the chain rule for derivatives, we find

$$\begin{aligned} \xi_1 &\simeq -3.9304 \times 10^{-3} \Delta \tilde{P}_\phi^e \\ \xi_2 &\simeq 3.3571 \times 10^{-3} \Delta \tilde{\lambda}^e \end{aligned} \quad (80)$$

From which we obtain

$$\begin{aligned}\Delta\hat{P}_\phi^e &= 182.278 & ; & \quad \Delta\hat{\lambda}_0^e \rightarrow \infty & ; & \quad \Delta\hat{\lambda}_1^e = 581.268 \\ (\Delta\hat{P}_\phi^i)^{-2} &\sim \mathcal{O}(\epsilon^2) & ; & \quad \Delta\hat{\lambda}_0^i \rightarrow \infty & ; & \quad \Delta\hat{\lambda}_1^i = 286.236\end{aligned}\tag{81}$$

Hence, the electron and ion density distribution functions finally read

$$\begin{aligned}\mathcal{F}^{e0} &\propto \exp\left[-\left(\frac{\hat{w} - 0.5105}{0.3096}\right)^2 - \left(\frac{\hat{P} - 0.1651}{182.278}\right)^2 - \left(\frac{\hat{\lambda} - 0.5246}{581.268}\right)^2\right] |\mathcal{J}| \\ \mathcal{F}^{i0} &\propto \exp\left[-\left(\frac{10^3 \times \hat{w} - 0.2780}{0.1686}\right)^2 - \left(\frac{\hat{\lambda} - 0.3984}{286.236}\right)^2\right] |\mathcal{J}|\end{aligned}\tag{82}$$

where the dimensionless variable $\hat{w} \equiv w/v_{the}^2$, with v_{the} computed at the center of the tokamak, has been introduced. Therefore, Eqs. (68) ensure that for collisional tokamak-plasmas in the Onsager region, the DDF $\mathcal{F}^0$ *coincides exactly* with the reference DDF estimated by the neoclassical theory. In the coordinates $\hat{w}$, $\hat{P}_\phi$ and $\hat{\lambda}$ (and ψ), the expression of the DDF is given by Eq. (19). Notice that in this case $c_2^{(1)} = 0$ ($\Delta\hat{\lambda}_0^\alpha \rightarrow \infty$) and the presence of the parameter $c_2^{(0)}$ (or of the parameter $\Delta\hat{\lambda}_1^\alpha$) is crucial.

V. CONCLUSIONS

Using statistical thermodynamics approach we have derived the general expression of the (density of) distribution function $\mathcal{F}^0$ for the case of a thermodynamic system out of equilibrium, subject to three thermodynamic forces. The local equilibrium is fixed by imposing the following conditions :

- i) The minimum entropy production condition on the two Prigogine's fluctuations α_1 and α_2 ;
- ii) The maximum entropy principle on the variable w , for $\alpha_1 = \alpha_2 = 0$;
- iii) The scale invariance of the restrictions used in the maximization of the entropy;
- iv) A new mathematical ansatz, used in selecting a minimal number of restrictions and implicitly free parameters.

From this ansatz results a singularity of the DDF that has immediate physical interpretation in terms of the intermittency in turbulent plasmas.

The derived DDF, $\mathcal{F}^0$, is more general than that currently used for fitting the numerical steady-state solution obtained by simulating ICRH plasmas and for describing various scenarios of tokamak plasmas. The adopted procedure can be generalized for systems subjected to an arbitrary number of thermodynamic forces. By kinetic theory, we have linked, and then fixed, the seven free parameters entering in $\mathcal{F}^0$ with the external energy sources and the (internal) entropy production source strength. To be more concrete, we have analyzed the case of, fully ionized, magnetically confined plasmas.

A perspective of this work is the following. In 2009, a macroscopic theory for closure relations has been proposed for systems out of Onsager's region [28]. This theory is referred to as the thermodynamic field theory (TFT). The aim of the work is to determine the nonlinear flux-force relations that respect the thermodynamic theorems for systems far from equilibrium. Through the TFT it is possible to estimate the nonlinear DDFs for collisional tokamak-plasmas when the nonlinear contributions cannot be neglected [29]. Also in this case, the crucial question is : *When does our reference DDF identify with the, steady-state, nonlinear distribution function derived by the TFT ?* The solution of this difficult problem will contribute to establish the link between the macroscopic approach (TFT) with a microscopic description based on statistical thermodynamics. Another problem to be solved is the possibility to improve the numerical fit by adding new free parameters according to the principles exposed in the Appendix. The work is ongoing.

VI. ACKNOWLEDGMENTS

Dr. Fulvio Zonca, of the Association EURATOM-ENEA, in Frascati (Italy), inspired the realization of this work. We are indebted to him for his invaluable support and his scientific suggestions. One of us (G. Sonnino), is very grateful to Prof. M.Malek Mansour, of the Université Libre de Bruxelles, for his scientific suggestions and for his help in the development of this work. M.Tlidi is a Research Associate with the Fonds de la Recherche Scientifique F.R.S.-FNRS, Belgium.

Appendix A: The Physical Justification of the Gamma Distribution Function

In order to justify the w dependence of $\mathcal{P}(w)$ we will use the principle of maximal entropy with suitable chosen restrictions obtained from scale invariance and a new mathematical ansatz. We denote by $S[\rho(.)]$ the entropy functional of a given probability density function $\rho(w)$:

$$S[\rho(.)] = - \int_0^\infty \rho(w) \log(\rho(w)) dw \quad (\text{A1})$$

The maximum entropy principle will be applied by imposing a set of restrictions:

$$\int_0^\infty \rho(w) dw = 1 \quad (\text{A2})$$

$$\int_0^\infty w \rho(w) dw = E(w) = \mu_1 \quad (\text{A3})$$

$$\int_0^\infty \log(w) \rho(w) dw = E(\log(w)) = \nu \quad (\text{A4})$$

The naturalness of the restriction given by Eq. (A3) is clear: we observe that $\mu_1 = T/m$ where T is the temperature. Concerning the naturalness of the choice of restriction (A4), we remark that all of the previous restrictions belongs to a class that is invariant under scaling. More generally, for a restriction of the form $E(w^\alpha) = \mu_\alpha$, respectively for $E(\log(w)) = \nu$, the effect scaling $w = kw'$ is $E(w'^\alpha) = \mu'_\alpha = \mu_\alpha k^{-\alpha}$, respectively for $E(\log(w')) = \nu' = \nu - \log(k)$. The problem why the PDF obtained by the one of the possible simplest choice from the more general set of restrictions

$$E(w^{\alpha_k}) = \mu_k ; \quad \text{with} \quad \mu_0 \equiv 1; \alpha_0 = 0; \alpha_1 = 1; k = 0, 1, 2, \dots, \quad (\text{A5})$$

$$E(\log(w)) = \nu \quad (\text{A6})$$

deserves further study. A partial answer is given by simplicity and extremality reasons. Observe first that the restriction (A6) can be seen as a limiting case of (A5). Indeed, suppose that we have for some fixed k : $\alpha_k = \varepsilon \ll 1$

$$E(w^\varepsilon) = \mu_k \quad (\text{A7})$$

From (A2) and (A7) results

$$E\left(\frac{w^\varepsilon - w^0}{\varepsilon}\right) = \frac{\mu_k - 1}{\varepsilon} \quad (\text{A8})$$

When the support of the PDF $\rho(w)$ is concentrated mainly on the domain when $|\log(w)|$ is not too large, then we can approximate: $(w^\varepsilon - w^0)/\varepsilon \cong \log(w)$, so the Eq.(A8) is

reduced to Eq.(A4). So, the restriction of the type (A4) can be seen as the an extreme case. It is easily checked that applying the maximal entropy principle from A6 results $\rho(w) \underset{w \rightarrow 0}{\asymp} \text{const } w^{\gamma-1}$ for some $\gamma > 0$. Notice that when $\gamma < 1$, the singularity of our DDF is related to the *intermittency* shown by real physical DDFs [30], with time and ensemble average are provided by our DDF. On the other hand, it is easily checked that in the case of restrictions, A5 after applying the maximal entropy principle, we obtain $\rho(w) \underset{w \rightarrow \infty}{\asymp} c_1 \exp(-c_2 w^m)$ for some $c_1, c_2 > 0$, where $m = \max\{\alpha_1, \dots, \alpha_n\}$. But $m > 1$ give a much faster decay compared to Maxwell distribution, so we obtain $m < 1$. *Consequently, we use the minimal ansatz given by Eqs (A2)-(A4), for the selection of the restrictions.* From this reason of extremality of the restriction Eq.(A4) and motivated by the minimality assumptions, we will explore the consequences of the minimal and extremal model given by Eqs (A2)-(A4). In order to find the distribution function $\rho(w)$ that maximizes the entropy (A1), with restrictions (A2)-(A4) we use the Lagrange multiplier method. The result is the gamma distribution function, given by

$$\rho(w) = \frac{\Theta}{\Gamma(\gamma)} (w/\Theta)^{\gamma-1} \exp(-w/\Theta) \quad (\text{A9})$$

where $\Gamma(\gamma)$ is the Euler gamma function. The relation between the parameters γ , Θ and $\mu_1 = T/m$, ν is given by

$$E(w) = T/m = \gamma\Theta \quad (\text{A10})$$

$$E(\log(w)) = \nu = \Psi(\gamma) + \log(\Theta) \quad (\text{A11})$$

where $\Psi(\gamma)$ is the digamma function. Due to the very special mathematical peculiarity in the ansatz Eqs (A2)-(A4), the resulting PDF is expected to have also some special properties. Indeed, the gamma distribution is *infinitely divisible and stable* : if the PDF of independent random variables $X_1, \dots, X_n$ is a gamma distribution with the same scale parameter and shape parameters $\gamma_1, \dots, \gamma_n$, then PDF of the random variable $\sum_{k=1}^n X_k$ is again a gamma distribution with the same scale parameter and with shape parameter $\sum_{k=1}^n \gamma_k$. We are convinced that the physical interpretation of the infinite divisibility property is a challenging

problem.

-
- [1] R. Balescu, 1977 *Statistical Dynamics. Matter out of Equilibrium*, Imperial College Press, Printed in Singapore by Uto-print.
 - [2] R.F. Greene and H.B. Callen, *Phys. Rev.*, **83**, 231, (1951).
 - [3] R.C. Tolman, 1938 *The Principle of Statistical Mechanics*, Oxford University Press, New York.
 - [4] R.H. Fowler, 1936 *Statistical Mechanics*, Cambridge University Press, New York.
 - [5] I. Prigogine, 1954 *Thermodynamics of Irreversible processes*, (John Wiley & Sons).
 - [6] R. Balescu, 1988 *Transport Processes in Plasmas. Vol 1. Classical Transport*, Elsevier Science Publishers B.V., Amsterdam, North-Holland.
 - [7] S.R. De Groot and P. Mazur, 1984 *Non-Equilibrium Thermodynamics*, Dover Publications, Inc., New York.
 - [8] G. Sonnino *Phys.Rev. E*, **79**, 051126, (2009).
 - [9] A. Cardinali *et al.*, *23rd IAEA Fusion Energy Conference*, (2010).
 - [10] A. Pizzuto *et al.*, *Nucl. Fusion*, **50**, 95005, (2010).
 - [11] F. Zonca and L. Chen, *Phys. Plasmas*, **7**, 4600, (2000).
 - [12] R. Balescu, 1988 *Transport Processes in Plasmas. Vol 2. Neoclassical Transport*, Elsevier Science Publishers B.V., Amsterdam, North-Holland.
 - [13] I. Prigogine (1947), *Etude Thermodynamique des Phénomènes Irréversibles*, Thèse d'Aggrégation de l'Enseignement Supérieur de l'Université Libre de Bruxelles (U.L.B.).
 - [14] Y. Park, Sung, K. Bera, Anil, *Journal of Econometrics* (Elsevier), 219 (2009). Retrieved (2011).
 - [15] A. Papoulis and S.U. Pillai, 2002 *Probability, Random Variables and Stochastic Processes*, Mac Graw Hill, Fourth edition.
 - [16] J. Wesson, 2004 *Tokamaks*, Oxford Science Publications, Clarendon Press, Oxford.
 - [17] W.A. Cooper *et al.*, *Nucl. Fusion*, **46**, 683, (2006).
 - [18] J. Freidberg, 2007 *Plasma Physics and Fusion Energy*, Cambridge Press University, Cambridge.
 - [19] F.L. Hinton and M.N. Rosenbluth, *Physics Letters A*, **259**, 267, (1999).
 - [20] C. F.F. Karney, *Comp. Phys. Report*, **4**, 183 (1986).

- [21] M. Brambilla, *Nuclear Fusion*, **34**, No.8, 1121 (1994).
- [22] A. Cardinali, S. Briguglio, G. Calabrò, *et Al.*, *Nuclear Fusion*, **49**, 095020 (2009).
- [23] C. W. Gardiner, 2004 *Handbook of Stochastic Methods: for Physics, Chemistry and the Natural Sciences*, Third edition Springer Verlag, Berlin, Heidelberg, New York.
- [24] W. Horsthemke and R. Lefever, 1985 *Noise Induced Transitions. Theory and Applications in Physics, Chemistry and Biology*, Springer Series in Synergetics, Vol. 15), Berlin, Heidelberg.
- [25] C. S. Chang and F. L. Hinton, *Phys. Fluids*, **25**, 1493 (1982).
- [26] F.L. Hinton and R.D. Hazeltine, *Rev. Mod. Phys.*, **48**, 239, (1976).
- [27] G. Sonnino and P. Peeters *Physics of Plasmas* **15**, 062309/1-062309/23 (2008).
- [28] G. Sonnino, *Phys. Rev. E.*, **79**, 051126 (5 Pt 1), (2009).
- [29] G. Sonnino, *The European Physical Journal D (EPJD)*, **62**, Issue 1, 81, (2011).
- [30] S. Aumaitre, F. Petrelis, K. Mallick, *Phys. Rev.Lett.* **95**, Issue 6, 064101, (2005).
- [31] We would like to stress that, strictly speaking, the entropy production reaches an extremum when its first partial derivatives with respect to the three fluctuations, α_1, α_2 and α_3 , vanish at $\alpha_1 = \alpha_2 = \alpha_3 = 0$. In this paper, we shall characterize the *equilibrium state* as the state where the first two partial derivatives of the entropy production, made with respect to the two fluctuations α_1 and α_2 , vanish at $\alpha_1 = \alpha_2 = 0$. α_3 is, however, interpreted as a random variable satisfying, at the equilibrium state, the *maximum entropy principle* (see later in this section). This is the reason of why, in our assumption, we use the expression "the entropy production *tends* to reach $\dots$ " and not "the entropy production *reaches* $\dots$ ".
- [32] We recall that, by definition, the degrees of advancement variables ζ_j satisfy the condition $\lim_{\zeta_j \rightarrow \zeta_j^0} \xi_i = 0$ [5].
- [33] Of course, Eqs. (9) may be easily generalized. This would lead to a final expression for the distribution function, which is even more general than Eq. (19). However, to avoid complicating further the formalism, we preferred to work with the simplest case provided by Eqs (9).
- [34] Here, we shall limit ourselves to analyze the so-called *simplified, quasi-linear, Fokker Planck equation* (QLFPE) where it is taken into account only the contribution due to the perpendicular component of the electric field, $E_{\perp}^+$, which is concordant to the direction of rotation of the minority [21]. Eq. (??) neglects then the contributions due to the perpendicular component of the electric field, $E_{\perp}^-$, which is discordant to the direction of rotation of the minority, and the one due to the parallel component of the electric field, $E_{\parallel}$.

- [35] Note that in the limit $\hat{\eta} \equiv r/R_0 \ll 1$, the ratio between the poloidal component and the toroidal component of the magnetic field strength is independent of the poloidal angle θ (at the first approximation in $\hat{\eta}$).
- [36] In our calculations, we shall assume that, other than $v_{\parallel}$, also $dv_{\parallel}$ is close to 0.
- [37] At the values $X_e^1 = X_e^2 = 0$ and $X_i^1 = 0$, $\mathcal{F}_{\alpha}^0$ exactly coincides with a gamma distribution function. Since the entropy productions, Eqs (52), have been estimated by truncating the expressions at the second order in the drift parameters ϵ [6], for coherence, Eqs (76) and (77) should also be solved up to the second order in ϵ . We are then imposing the conditions for which $\mathcal{F}_{\alpha}^0$ approximates the gamma distribution function up to ϵ^2 .